

Drug Information Sheet("Kusuri-no-Shiori")

Internal

Published: 04/2016

The information on this sheet is based on approvals granted by the Japanese regulatory authority. Approval details may vary by country. Medicines have adverse reactions (risks) as well as efficacies (benefits). It is important to minimize adverse reactions and maximize efficacy. To obtain a better therapeutic response, patients should understand their medication and cooperate with the treatment.

Brand name: Selbex Capsules 50mg

Active ingredient: Teprenone

Dosage form: grayish blue-green/light orange capsule, whole length 14.3 mm

Print on wrapping: (Face) セルベックス 50mg, SX50E

(Back) セルベックス50mg, SX50E, Selbex, 50mg, セルベックス

**Effects of this medicine**

This medicine increases secretion of gastric mucus, and promotes protection and repair of damaged gastric mucosa. It is usually used in the treatment of gastritis and gastric ulcer.

Before using this medicine, be sure to tell your doctor and pharmacist

- If you have previously experienced any allergic reactions (itch, rash, etc.) to any medicines.
- If you are pregnant or breastfeeding.
- If you are taking any other medicinal products. (Some medicines may interact to enhance or diminish medicinal effects. Beware of over-the-counter medicines and dietary supplements as well as other prescription medicines.)

Dosing schedule (How to take this medicine)

- Your dosing schedule prescribed by your doctor is<< to be written by a healthcare professional>>
- In general, for adults, take 1 capsule (50 mg of the active ingredient) at a time, 3 times daily, after meal. The dosage may be adjusted according to the disease, age or symptoms. Strictly follow the instructions.
- If you miss a dose, take a dose as soon as possible when you remember that you missed a dose. However, if it is almost time for the next dose, skip the missed dose and continue your regular dosing schedule. You should not take two doses at one time.
- If you accidentally take more than your prescribed dose, consult with your doctor or pharmacist.
- Do not stop taking this medicine unless your doctor instructs you to do so.

Precautions while taking this medicine**Possible adverse reactions to this medicine**

The most commonly reported adverse reactions include rash, itching, constipation, diarrhea, thirst, abdominal pain, feeling of enlarged abdomen and headache. If any of these symptoms occur, consult with your doctor or pharmacist.

The symptoms described below are rarely seen as initial symptoms of the adverse reactions indicated in brackets. If any of these symptoms occur, stop taking this medicine and see your doctor immediately.

- nausea, vomiting, loss of appetite, yellowing in the skin or conjunctiva ocular [hepatic function disorders, jaundice]

The above symptoms do not describe all the adverse reactions to this medicine. Consult with your doctor or pharmacist if you notice any symptoms of concern other than those listed above.

Storage conditions and other information

- Keep the medicine out of the reach of children. Store it away from direct sunlight, heat and moisture.
- Do not store the remainder and discard it.

For healthcare professional use only / /

For further information, talk to your doctor or pharmacist.